

Case 3028***Aphanius* Nardo, 1827 (Osteichthyes, Cyprinodontiformes): proposed placement on the Official List**

Maurice Kottelat

Route de la Baroche 12, Case postale 57, 2952 Cornol, Switzerland (address for correspondence); Department of Zoology, National University of Singapore, Kent Ridge, Singapore 119260 (e-mail: mkottelat@dplanet.ch)

Alwyne Wheeler

clo Department of Zoology, The Natural History Museum, Cromwell Road, London SW7 5BD, U.K.

Abstract. The purpose of this application is to conserve the name *Aphanius* Nardo, 1827 for a genus of Palearctic fishes (family CYPRINODONTIDAE). The name has been in uninterrupted use since at least 1926 but a few authors have recently replaced it with *Lebias* Goldfuss, 1820, a name which, with a single exception in 1895, had remained unused since 1846 until resurrected in 1995 and which does not refer to the same taxon as *Aphanius*. *Aphanius* includes at least 17 extant species and fossil remains have been reported from Miocene deposits.

Keywords. Nomenclature; taxonomy; *Aphanius*; *Cyprinodon*; *Lebias*; CYPRINODONTIDAE; *Aphanius fasciatus*; *Cyprinodon variegatus*; tooth carps; freshwater; brackish water; Palearctic.

1. The name *Aphanius* Nardo, 1827 (ref. 1827a, pp. 34, 39-40; also published in 1827b, col. 487) relates to a genus of fresh and brackish water tooth carps (family CYPRINODONTIDAE) with a wide distribution, basically peri-Mediterranean, extending from Portugal and Morocco to Pakistan. The genus was established with two originally included nominal species, *A. nanus* and *A. fasciatus*, both of Nardo (1827a, pp. 34, 40; 1827b, col. 488). Jordan (1917, p. 121) selected *A. nanus*, a junior synonym of *Lebias fasciata* Valenciennes in Cuvier & Valenciennes, 1827, as the type species (see para. 9). The genus now includes at least 17 species and subspecies, although it is likely that the number is much greater, and several have very restricted distributions in arid zones. Several populations and species are now seriously threatened by depletion of water resources for urban and agricultural use, pollution and introductions and are given local protection and listed by international agencies (for example, the 1996 IUCN Red List of Threatened Animals). Fossil remains identified as *Aphanius* have been reported from Miocene deposits.

2. The name *Aphanius* has been in uninterrupted use since at least 1926 but it has recently been treated by a few authors as a junior synonym of *Lebias* Goldfuss, 1820. This latter name has for more than 150 years been considered a junior synonym of *Cyprinodon* La Cépède, 1803 and, with a single exception in 1895, remained unused since 1846 until resurrected in 1995. However, the (1995) type species designation

which was supposed to render *Aphanius* a synonym of *Lebias* is invalid. To secure the continued and unhindered use of the name *Aphanius* we propose that it be placed on the Official List.

3. The history of the name *Lebias* is as follows. Cuvier (1816, p. 199) proposed a new genus of tooth carps using the vernacular name 'Le *Lebias*'. He did not mention species by name. Oken (1817, p. 1183), in a commentary on Cuvier's classification and a comparison between the latter and his own, adopted the Latin name *Lebia* and, by reference to Cuvier, rendered the name available (see Gill, 1903, p. 967). There were no included species. The name *Lebia* Oken is, however, a junior homonym of *Lebia* Latreille, 1802, a much used name in Coleoptera.

4. The tooth carp genus was subsequently briefly described by Goldfuss (1820, p. 16) who, like Oken, referred to Cuvier (1816) but used the spelling *Lebias*. Since *Lebia* Oken is a junior homonym (para. 3 above), the synonym *Lebias* Goldfuss, 1820 would be the valid name for the genus (if it were separated from *Cyprinodon* La Cépède, 1803; see para. 6 below). Again there were no originally included species (Goldfuss noted 'Arten sind noch nicht beschrieben'). The first subsequent mention of the genus, which also included a nominal species, was by Le Sueur (1821) who, like Goldfuss (1820), referred to Cuvier (1816) but used *Lebia*, the same spelling as Oken. Le Sueur (p. 6) placed in the genus the single nominal species *Lebia ellipsoidea* Le Sueur, 1821 from Florida. Also in 1821, Valenciennes (in Humboldt & Valenciennes, p. 159) referred to the genus 'que M. Cuvier a établi' under the name *Lebias* and named Cuvier's (1816) two new species: *L. rhomboidalis* Valenciennes, 1821 (p. 160, pl. 61, figs. 3, 7) from North America and *L. fasciata* Valenciennes, 1821 from Europe (p. 160, pl. 61, fig. 4). Le Sueur's work was published in January 1821 (as recorded in vol. 2 of the *Journal of the Academy of Natural Sciences of Philadelphia*, in vol. 1, p. 8 of the *Proceedings of the Academy* . . . (1841), and in the 'Index to the scientific contents of the Journal and Proceedings of the Academy . . . 1812–1912' (1913)), while Valenciennes's publication can be dated only to the year 1821 (see Sherborn, 1899, p. 428; Lazara, 1993, p. 1160; and Kottelat, 1997, p. 162). It is clear from Oken (1817), Goldfuss (1820) and other early authors that both *Lebia* and *Lebias* are spellings derived from Cuvier's (1816) vernacular 'Le *Lebias*' and no author (except Lazara, 1995; see para. 8 below) has ever regarded them as distinct. *Lebia ellipsoidea* Le Sueur, 1821 from Florida is thus the type species of *Lebias* Goldfuss, 1820 by subsequent monotypy.

5. Lazara (1995), putatively acting as the First Revisor, selected *Lebia* as the valid spelling from Le Sueur (1821). His action was invalid, however, because both the spellings *Lebia* and *Lebias* had been published before Le Sueur (1821), by Oken (1817) and Goldfuss (1820) respectively. In any case, Le Sueur (pp. 2, 5, 7) consistently adopted the spelling *Lebia*; on p. 5 'the *Lebias*' was a plural vernacular use, and on pl. 2 the spelling '*Lebias*' was probably an engraver's error (the specific name *ellipsoidea* was misspelt as '*elipsoides*', and the generic name '*Mollinesia*' in the text was spelt '*Molienisia*' on pl. 3, both being misspellings of *Mollienesia*).

6. In 1846 Valenciennes (in Cuvier & Valenciennes, p. 145) included both the New and Old World cyprinodont species in *Cyprinodon* La Cépède, 1803, giving *Lebias* (which he cited from Cuvier, 1816) as a junior synonym. Valenciennes synonymised his (1821) American species *L. rhomboidalis* with *C. variegatus* La Cépède, 1803 (pp. 486, 487), described from Charleston Bay and the type species of *Cyprinodon* by

monotypy. Valenciennes (pp. 146–151) considered that Cuvier (1816) had made a number of errors in his original description of ‘Lebias’, and stated that this description and those of the two nominal species *C. variegatus* and *L. rhomboidalis* had all been based on the same two specimens in the Muséum National d’Histoire Naturelle in Paris. In discussing Cuvier’s work he noted ‘Il y a là une suite de méprises; car il est évident que le genre *Lebias* a été créé pour un poisson qui n’est autre chose que le cyprinodon varié’ (i.e. *Cyprinodon variegatus*). Valenciennes (1846, pp. 173–178) also included in *C. variegatus* the nominal species *Lebias* (sic) *ellipsoidea* Le Sueur, 1821, new material from Lake Pontchartrain, near New Orleans, having been sent to the Paris Museum. Valenciennes (1846, pp. 156–159) retained the name *C. fasciatus* (Valenciennes, 1821) for the European cyprinodont species. Günther (1866, pp. 302, 305) also listed New and Old World cyprinodont species under *Cyprinodon* La Cepède, 1803, citing *Lebias* and *Aphanius* as synonyms and, like Valenciennes (1846), considered *C. variegatus*, *L. rhomboidalis* and *L. ellipsoidea* to refer to the same species. The synonymy of *L. ellipsoidea* with *C. variegatus* rendered the name *Lebias* Goldfuss, 1820 a junior subjective synonym of *Cyprinodon*.

7. Garman (1895, p. 20) also cited *L. ellipsoidea* Le Sueur, 1821 as a synonym of *C. variegatus* La Cepède, 1803; he used (pp. 19–29) the name *Cyprinodon* for New World species and (pp. 29–34), overlooking the consequences of the synonymy of *L. ellipsoidea* with *C. variegatus*, resurrected *Lebias* for Old World species, including *C. fasciatus* (Valenciennes, 1821), and treated *Aphanius* as a junior synonym. Like Günther (1866; see para. 6 above), Boulenger (1907, pp. 406–412) used *Cyprinodon* for both New and Old World species, citing *Lebias* and *Aphanius* as synonyms, and the name *Lebias* dropped from use. Hubbs (1926, p. 16) again separated New and Old World species, adopting the names *Cyprinodon* and *Aphanius* respectively. He was followed by Myers (1931), who commented (p. 12) that ‘*Lebias* is a synonym of *Cyprinodon*, and the European forms belong to *Aphanius*’, Myers (1935, p. 303) and Miller (1948, p. 21), who commented that *Aphanius* was ‘formerly [i.e. by Garman, 1895] called *Lebia* or *Lebias*, a synonym of *Cyprinodon*’. The name *Lebias* had not been used for more than a century until resurrected by Lazara in 1995.

8. Lazara (1995) attempted to separate the spellings *Lebia* and *Lebias* and to apply them to different taxa. He recognised *Lebia*, as of Le Sueur (1821), as a junior synonym of *Cyprinodon*, and by designating *Lebias fasciata* Valenciennes, 1821 as the type species of *Lebias* Goldfuss, 1820, sought to reintroduce *Lebias* in place of *Aphanius* Nardo, 1827 as the name for Old World cyprinodonts (see para. 9 below). As recorded in para. 4 above, *Lebias* and *Lebia* are variant spellings of the same name, i.e. both were based on Cuvier’s ‘Le Lebias’, and *Lebias*, dating from Goldfuss (1820), is the (potentially) valid spelling. The type species of *Lebias* is the American species *Lebia ellipsoidea* Le Sueur, 1821 by subsequent monotypy and Lazara’s (1995) type species designation is invalid. Lazara (1995) acknowledged that the name *Aphanius* had been in use for many years.

9. As noted in para. 1 above, Jordan (1917) selected the first of the nominal species (*A. nanus* Nardo, 1827) included in *Aphanius* Nardo, 1827 as the type species of the genus. He recorded *Aphanius* as a valid genus ‘replacing *Lebias* of authors (not of Cuvier)’. *Aphanius nanus* has been treated as a synonym of *A. fasciatus* Nardo, 1827 and of *Lebias fasciata* Valenciennes, 1821 since at least Garman (1895, pp. 29, 30)

and Boulenger (1907, p. 407), and *A. fasciatus* (Valenciennes, 1821) is thus the valid name for the type species of *Aphanius* (see Wildekamp, 1993, p. 48).

10. Very few authors have followed Lazara (1995) in his use of the name *Lebias* in place of *Aphanius*. The overwhelming use is of the latter name, both in works on taxonomy of Recent and fossil species and in the applied fields of biology, reproduction, genetics, biochemistry, hybridisation, physiology and ecology. Recent representative works, covering systematics, checklists, field guides and conservation documents, in which *Aphanius* has been used are Economidis (1991, 1992), Doadrio, Elvira & Bernat (1991), Gandolfi, Zerunian, Torricelli & Marconato (1991), Wildekamp (1993), Coad (1996), Ferrito & Tigano (1996), Maitland & Crivelli (1996), Kottelat (1997), Maitland (2000), and several papers in the publications edited by Crivelli & Maitland (1995) and by Kirchhofer & Hefti (1996). A search of *Zoological Record on CD* (vols. 115–136) showed a further 162 publications in which the name *Aphanius* has been used between 1978 and 2000 (the complete list is held by the Secretariat of the Commission). It is very desirable that the use of the name *Aphanius* be continued in local, regional, national and international legal instruments, conservation policy documents and Red Lists; a change of name would seriously threaten the efficiency of conservation measures for many of the species concerned, several of which are in danger of immediate extinction.

11. As demonstrated above, the names *Lebia* Oken, 1817 and *Lebias* Goldfuss, 1820 both refer to the same taxon, as had always been accepted until Lazara (1995). The type species is *Lebia ellipsoidea* Le Sueur, 1821 (see para. 4 above), which is a junior subjective synonym of *Cyprinodon variegatus* La Cepède, 1803, the type species of *Cyprinodon* La Cepède, 1803; accordingly *Lebia* and *Lebias* are junior synonyms of *Cyprinodon*. However, Lazara (1995), following Garman (1895), misinterpreted *Lebias* and adopted it instead of *Aphanius* Nardo, 1827 as the valid name for Old World species of tooth carps, even though he acknowledged that *Aphanius* had been in use for many years. In the interests of stability and to avoid misunderstanding, we propose that *Aphanius* should be placed on the Official List and that the name *Lebias* Goldfuss, 1820 should be suppressed.

12. The International Commission on Zoological Nomenclature is accordingly asked:

- (1) to use its plenary power to suppress the name *Lebias* Goldfuss, 1820 for the purposes of the Principle of Priority but not for those of the Principle of Homonymy;
- (2) to place on the Official List of Generic Names in Zoology the name *Aphanius* Nardo, 1827 (gender: masculine), type species by subsequent designation by Jordan (1917) *Aphanius nanus* Nardo, 1827 (a junior subjective synonym of *Lebias fasciata* Valenciennes in Humboldt & Valenciennes, 1821);
- (3) to place on the Official List of Specific Names in Zoology the name *fasciata* Valenciennes in Humboldt & Valenciennes, 1821, as published in the binomen *Lebias fasciata* (senior subjective synonym of the specific name of *Aphanius nanus* Nardo, 1827, the type species of *Aphanius* Nardo, 1827);
- (4) to place on the Official Index of Rejected and Invalid Generic Names in Zoology the following names:
 - (a) *Lebia* Oken, 1817 (a junior homonym of *Lebia* Latreille, 1802);
 - (b) *Lebias* Goldfuss, 1820 (suppressed in (1) above).

References

- Boulenger, G.A.** 1907. *Zoology of Egypt. The fishes of the Nile*. li, 578 pp. 97 pls. Rees, London.
- Coad, B.W.** 1996. Systematics of the tooth-carp genus *Aphanius* Nardo, 1827 (Actinopterygii: Cyprinodontidae) in Fars Province, southern Iran. *Biologia* (Bratislava), **51**(2): 163–172.
- Crivelli, A.J. & Maitland, P.S.** (Eds.). 1995. Endemic freshwater fishes of the northern Mediterranean region. *Biological Conservation*, **72**: 121–337.
- Cuvier, G.L.** 1816. *Le Règne Animal distribué d'après son organisation . . . Les reptiles, les poissons, les mollusques et les annélides*, vol. 2. Déterville, Paris.
- Doadrio, I., Elvira, B. & Bernat, Y.** 1991. *Peces continentales españolas. Inventario y clasificación de zonas fluviales*. 221 pp. Icona. Madrid.
- Economidis, P.S.** 1991. *Check list of freshwater fishes of Greece*. 48 pp. Hellenic Society for the Protection of Nature, Athens.
- Economidis, P.S.** 1992. Fishes. Pp. 43–81 in Karandinos, M. & Paraschi, L. (Eds.), *The Red Data Book of threatened vertebrates of Greece*. Hellenic Zoological Society & Hellenic Ornithological Society, Athens.
- Ferrito, V. & Tigano, C.** 1996. Decline of *Aphanius fasciatus* (Cyprinodontidae) and *Salaria fluviatilis* (Blenniidae) populations in freshwaters of eastern Sicily. *Ichthyological Exploration of Freshwaters*, **7**(2): 181–184.
- Gandolfi, G., Zerunian, S., Torricelli, P. & Marconato, A.** 1991. *I pesci delle acque interne italiane*. 597 pp., 9 pls. Istituto Poligrafico e Zecca dello Stato, Roma.
- Garman, S.** 1895. The cyprinodonts. *Memoirs of the Museum of Comparative Zoology*, Harvard, **19**: 1–179.
- Gill, T.** 1903. On some fish genera of the first edition of Cuvier's *Règne Animal* and Oken's names. *Proceedings of the United States National Museum*, **26**: 965–967.
- Goldfuss, G.A.** 1820. *Handbuch der Zoologie*, vol. 2. xxiv, 510 pp., 4 pls. Schrag, Nürnberg.
- Günther, A.** 1866. *Catalogue of the fishes in the British Museum*. vol. 6. xv, 368 pp. British Museum, London.
- Hubbs, C.L.** 1926. Studies of the fishes of the Order Cyprinodontes. VI. Material for a revision of the American genera and species. *Miscellaneous Publications. Museum of Zoology, University of Michigan*, **16**: 1–87.
- Humboldt, A. de & Valenciennes, A.** 1821. Recherches sur les poissons fluviatiles de l'Amérique équinoxiale. Pp. 145–216, pls. 45–51 in Humboldt, A. de & Bonpland, A., *Voyage aux régions équinoxiales du nouveau continent, fait en 1799, 1800, 1801, 1802, 1803 et 1804*, vol. 2, part II (Recueil d'observations de zoologie et d'anatomie comparées faites dans l'Océan Atlantique, dans l'intérieur du nouveau continent et dans la mer du sud . . .). Schoell, Paris.
- IUCN.** 1996. *1996 IUCN Red List of threatened animals*. IUCN, Gland & Cambridge.
- Jordan, D.S.** 1917. The genera of fishes, from Linnaeus to Cuvier, 1758–1833, seventy-five years with the accepted type of each. A contribution to the stability of scientific nomenclature. *Leland Stanford Junior University Publications, University Series*, **27**: 1–161.
- Kirchhofer, A. & Hefti, D.** (Eds.) 1996. *Conservation of endangered freshwater fish in Europe*. xii, 341 pp. Birkhäuser, Basel.
- Kottelat, M.** 1997. European freshwater fishes. An heuristic checklist of the freshwater fishes of Europe (exclusive of former USSR), with an introduction for non-systematists and comments on nomenclature and conservation. *Biologia* (Bratislava), Section Zoology, **52** (Suppl. 5): 1–271
- La Cèpède, B.** 1803. *Histoire naturelle des poissons*, vol. 5. xlviii, 803 pp., 21 pls. Paris.
- Lazara, K.J.** 1993. Dating and authorship of new species described in Humboldt and Valenciennes' 'Recherches sur les poissons fluviatiles de l'Amérique équinoxiale'. *Copeia*, **1993**(4): 1160–1161.
- Lazara, K.J.** 1995. History of the genera *Lebia* Oken 1817 and *Lebias* Goldfuss 1820 (Teleostei: Cyprinodontiformes: Cyprinodontidae) with designation of a type species for *Lebias*. *Copeia*, **1995**(2): 501–503.

- Le Sueur, C.A.** 1821. Description of a new genus, and of several new species of freshwater fish, indigenous to the United States. *Journal of the Academy of Natural Sciences of Philadelphia*, **2**: 2–8.
- Maitland, P.S.** 2000. *Guide to freshwater fish of Britain and Europe*. 256 pp. Hamlyn, London.
- Maitland, P.S. & Crivelli, A.J.** 1996. *Conservation des poissons d'eau douce*. 94 pp. Tour du Valat, Arles.
- Miller, R.R.** 1948. The cyprinodont fishes of the Death Valley system of eastern California and southwestern Nevada. *Miscellaneous Publications. Museum of Zoology, University of Michigan*, **68**: 1–155.
- Myers, G.S.** 1931. The primary groups of oviparous cyprinodont fishes. *Stanford University Publications, University Series, Biological Sciences*, **6**(3): 1–14.
- Myers, G.S.** 1935. An annotated list of the cyprinodont fishes of Hispaniola, with descriptions of two new species. *Zoologica*, **10**(3): 301–316.
- Nardo, J.D.** 1827a. Prodomus observationum et disquisitionum Adriaticae ichthyologiae. *Giornale di Fisica, Chimica, Storia naturale, Medicina ed Arti* (Pavia), Decade Seconde, **10**: 22–40.
- Nardo, J.D.** 1827b. Prodomus observationum et disquisitionum ichthyologiae Adriaticae. *Isis* (von Oken), **20**(6): 472–489.
- Oken, L.** 1817. Cuviers und Okens Zoologien neben einander gestellt. *Isis*, **1**: 1145–1185 (irregularly numbered).
- Sherborn, C.D.** 1899. A note on the date of the parts of 'Humboldt and Bonpland's Voyage; Observations de Zoologie'. *The Annals and Magazine of Natural History*, (7)**3**(17): 428.
- Valenciennes, A.** 1846. Vol. 18 in Cuvier, G.L. & Valenciennes, A., *Histoire naturelle des poissons*. xix, 505 pp. Paris.
- Wildekamp, R.H.** 1993. *A world of killies. Atlas of the oviparous cyprinodontiform fishes of the world*, vol. 1. 311 pp. American Killifish Association, Mishawaka.

Comments on this case are invited for publication (subject to editing) in the *Bulletin*; they should be sent to the Executive Secretary, I.C.Z.N., c/o The Natural History Museum, Cromwell Road, London SW7 5BD, U.K. (e-mail: iczn@nhm.ac.uk).